





Scanning Probe Microscopy on Active Polymer Solar Cells

David S. Ginger
Dept. of Chemistry
University of Washington
Seattle, WA 98195-1700

NIST Nov. 7, 2008



Acknowledgements

Group Members:

David Coffey, Ph.D. (physics)

Yeechi Chen

Brad MacLeod

Andrea Munro

Obadiah Reid

Joseph Wei

Keiko Munechika

Kevin Noone


Andreas Tillack

Deanna Rodovsky

Dr. Liam Pingree

Dr. Abhishek Kulkarni

Prof. Lee Park (Williams College)



Undergrads Co-Authoring Papers:

I-Kang Ding (to Stanford PhD)

Ilan Jen-La Plante (to Berkeley PhD)

Jessica Smith (to Berkeley PhD)

Marsha Ng (to UW MS)

Collaborators:

Glenn Bartholomew, Alex Jen, Younan Xia

Christine Luscombe, Mehmet Sarikaya, Dan Schwartz,

Francois Baneyx, Sam Jenekhe, Guozhong Cao, Daniel

Gamelin, Lee Park (Williams College)

Funding:

ONR

Camille and Henry Dreyfus Foundation

Alfred P. Sloan Foundation

Research Corporation, Cottrell Scholarship

DOE BES & DOE Solar America Initiative

NSF NIRT

PECASE (AFOSR)



NSF CAREER (DMR)

AFOSR DURIP and BIC program


NSF NER (EECS)

NSF STC MDITR

NSF UW MRSEC/"GEMSEC"



Physical Chemistry of Organic Electronics



optical processes near metals



Y. Chen, K. Munechika, I. Jen-La Plante, A. M. Munro, S. E. Skrabalak, Y. Xia, D. S. Ginger, *Appl. Phys. Lett.* **93** (2008). *in press*

L. Au, Y. Chen, F. Zhou, P. Camargo, Z.-Y. Li, D. S. Ginger, Y. Xia (2008) *in preparation*

Y. Chen & K. Munechika, D. S. Ginger, *MRS Bull.*, **33**(5) 536-542 (2008).


K. Munechika, J. M. Smith, Y. Chen, D. S. Ginger, *J. Phys. Chem. C*, **111**(51), 18906-18911 (2007).

Y. Chen, K. Munechika, and D. S. Ginger, *Nano Lett.*, **7**(3): 690-696 (2007).

B. J. Wiley, Y. Chen, J. McLellan, Y. Xiong, Z-Y. Li, D. S. Ginger, and Y. Xia, *Nano Lett.*, **7**(4): 1032-1036 (2007).

M. T. Zin, A. M. Munro, M. Gungormus, N.-Y. Wong, H. Ma, C. Tamerler, D. S. Ginger, M. Sarikaya, A. K.-Y. Jen, *J. Mater. Chem.*, **17**(9): 866-872, (2007).

quantum dot chromophores



A. M. Munro, D. S. Ginger, *Nano Lett.* (2008). (DOI: 10.1021/nl801132t)


A. M. Munro, J. A. Bardecker, M. S. Liu, Y. J. Cheng, Y. H. Niu, I. Jen La-Plante, A. K.-Y. Jen, D. S. Ginger, *Microchimica Acta* **160** (3): 345-350. (2008).

A. M. Munro, I. J.-L. Plante, M. S. Ng, D. S. Ginger, *J. Phys. Chem. C* **111**(17): 6220-6227 (2007).

Y. Niu, A. M. Munro, Y.-J. Cheng, Y. Tian, M. S. Liu, J. Zhao, J. A. Bardecker, I. J.-L. Plante, D. S. Ginger, A. K.-Y. Jen, *Adv. Mater.* **19**(20) 3371-3376 (2007).

J. Zhao, J. A. Bardecker, A. M. Munro, M. S. Liu, Y. Niu, I.-K. Ding, J. Luo, B. Chen, A. K.-Y. Jen, D. S. Ginger, *Nano Lett.* **6**(3), 463 (2006).

film morphology



L.Y. Park, A. M. Munro, D. S. Ginger, *J. Am. Chem. Soc (online ASAP)*

L. S. C. Pingree, O. G. Reid, D. S. Ginger *Adv. Mater.* (2008) *invited review In Press*

O. G. Reid, K. Munechika, D. S. Ginger *Nano Lett.* **8**(6): 1602-1609 (2008).

L. S. C. Pingree, B. A. MacLeod, D. S. Ginger *J. Phys. Chem C* **112** (21): 7922-7927 (2008).

L. S. C. Pingree, D. B. Rodovsky, D. C. Coffey, G. T. Bartholomew, D. S. Ginger *J. Am. Chem. Soc.* **129** (51): 15903 (2007)

D. C. Coffey, O. G. Reid, D. B. Rodovsky, G. T. Bartholomew, D. S. Ginger, *Nano Lett.* **7** (3): 738-744 (2007).


D. C. Coffey, D. S. Ginger, *Nature Materials* **5**(9): 735-740 (2006).

J. H. Wei, D. C. Coffey, D. S. Ginger, *J. Phys. Chem. B*, **110** (48): 24324-24330 (2006).

D. C. Coffey, D. S. Ginger, *J. Am. Chem. Soc.* **127**(13), 4564-4565 (2005).

Perspective on Scale

We need Really Big Solar Cells



Cell Area ~ 10^{11} m²

US Roadway:³ ~ 10^{10} - 10^{11} m²

rooftop capacity: 540 GW (18% of total energy)²
25% of highway area = 100% of electricity²



North American polyethylene capacity⁴ could cover Arizona with >4 layers of plastic wrap every year⁵
~300 days of newspaper

1: <http://www.ipcc.ch/pdf/assessment-report/ar4/wg3/ar4-wg3-chapter4.pdf>

2: http://www1.eere.energy.gov/solar/pdfs/set_myp_2007-2011_proof_1.pdf

3: <https://www.cia.gov/library/publications/the-world-factbook/print/us.html>

4: 30 billion pounds: <http://www.the-innovation-group.com/ChemProfiles/Polyethylene-HD.htm> Note: Polyethylene is NOT a semiconductor




5: given 0.5 mil thickness of "Glad wrap" = ~12.5 microns thick = <http://www.glad.com/faqs/plasticwrap.php>

6: circulation data USA today

Plastic Semiconductors are Printable




C&EN / Konarka



www.stanford.edu/group/mcgehee/research.html


ink jet printed flat panel TV




Ginger Lab, UW

Polymer Cells are Excitonic Solar Cells

film morphology critical to polymer photovoltaic performance





Morphology is a critical challenge



- 1) Can we develop tools to measure electronic properties (photocurrents, trapping rates, etc.)
- 2) Can we control film morphology (texture?)

Photoconductive AFM




D.C. Coffey, O.G. Reid, D.B. Rodovsky, G. P. Bartholomew, D.S. Ginger, *Nano Letters* 7 p738 (2007)

Small focus allows: 1) high intensity
2) less sample damage

MDMO-PPV/PCBM

Results: Photocurrent Variation



spin-coated from xylenes

Current


$$J_{SC}$$

$$FF$$

$$V_{OC}$$




all vary with location

Height



Variation in Chlorobenzene Blends


Current fluctuations on 2 length scales





$$\sigma_I = 0.57 \langle I \rangle$$

Good news: there is room to improve these devices with existing materials
However, source of large-scale variation unknown...suspect bottom contact
ITO, PEDOT?

Annealed P3HT/PCBM Blends



Looking at Local Charge Transport



O. G. Reid, K. Munechika, D. S. Ginger *Nano Lett.* 8(6): 1602-1609 (2008).


Mott-Gurney Law gives WRONG mobility in cAFM measurements (but is widely used!)

$$J = \frac{9}{8} \epsilon \epsilon_0 \mu \frac{V^2}{L^3}$$

Space Charge Limited Currents


$$J = \frac{9}{8} \epsilon \epsilon_0 \mu_0 e^{0.89 \gamma \sqrt{\frac{V}{L}}} \frac{V^2}{L^3}$$

cAFM and Mott-Gurney Law






$$J = \frac{9}{8} \epsilon \epsilon_0 \mu \frac{V^2}{L^3}$$

Clear deviation from
distance dependence
of Mott-Gurney law
in cAFM



O. G. Reid, K. Munechika, D. S. Ginger *Nano Lett.* 8(6): 1602-1609 (2008).

cAFM and Mott-Gurney Law






Numerically solve:

$$-\frac{d\rho}{dt} = \nabla \cdot (\mu\rho\nabla V + D\nabla\rho)$$

$$-\nabla^2 V = \frac{\rho}{\epsilon}$$


Geometry Accounts for Deviation



$$J_{c-AFM} = \alpha \mu_0 e^{0.89 \gamma \sqrt{\frac{V}{L}}} \epsilon \epsilon_0 \frac{V^2}{L^3} \delta \left(\frac{L}{d} \right)^{1.6 \pm 0.1}$$

O. G. Reid, K. Munechika, D. S. Ginger *Nano Lett.* 8(6): 1602-1609 (2008).

Electrostatic Force Microscopy (EFM)



An AC mode AFM technique:


$$\Delta\omega_{resonant} = -\frac{\omega_o}{2k} \frac{\partial Force}{\partial z}$$

$$Force = \frac{1}{2} \frac{\partial C}{\partial z} (V_{tip} - V_{surface})^2$$

Charge distribution
and
dielectric Surface Potential


$$\Delta Frequency = -\frac{\omega_o}{4k} \frac{\partial^2 C}{\partial z^2} (V_{tip} - V_{surface})^2$$

Carriers Change V_{surf} and d^2C/dz^2




$$\Delta \text{Frequency} = -\frac{\omega_o}{4k} \frac{\partial^2 C}{\partial z^2} (V_{\text{tip}} - V_{\text{surface}})^2$$

Where does the photocurrent come from?




“We observe a linear trend, giving substantial evidence that the external quantum efficiency scales with the interfacial area between the mesoscale phase separated regions.” Snaith et al. *Nano Lett.* p1353-1357 (2002)

“...measurements show that current is generated within the bulk and not at the boundaries of the micron-sized phase segregated features” McNeill et al., *Nano Lett.* p2503-2507 (2004)




“F8BT”



“PFB”

*Friend
Dastoor
Arias
Silva
Snaith
et al.*


More Efficient Blends Create Charge Faster



These are all *single-point* measurements: we can create a map!


Mapping Carrier Generation

Height




(spin-coated from xylene)

Charging Rate



EFM Measurements Predict Device Efficiency




What is happening as the films become more efficient?

D. C. Coffey and D. S. Ginger, *Nature Materials* 5, pp. 735-740, (2006)

trEFM on polyfluorenes summary

Where does the photocurrent come from?

“We observe a linear trend, giving substantial evidence that the external quantum efficiency scales with the interfacial area between the mesoscale phase separated regions.” Snaith et al. *Nano Lett.* p1353-1357 (2002)




PFB/F8BT

“...measurements show that current is generated within the bulk and not at the boundaries of the micron-sized phase segregated features” McNeill et al, *Nano Lett.* p2503-2507 (2004)



D. C. Coffey and D. S. Ginger, *Nature Materials* 5, pp. 735-740, (2006)

Local composition is most important in PF blends

trEFM is Superior to SKPM




*simple
SKPM
can give
wrong
answer*



unpublished






Can watch charges leak out of traps



(also decrease in total Δf)

Summary

- Scanning probe techniques help reveal microscopic workings of organic photodiodes
- Should think about polymer cells as consisting of many different nanoscale diodes in parallel
- Improved morphological control and anode/contact interface are promising targets for improving efficiencies with many materials
- Distribution of currents will be critical for design of multi-layer cells
- Surface chemistry promising for controlling lateral (and vertical?) morphology



ERROR: stackunderflow
OFFENDING COMMAND: ~

STACK: